LIVESTOCK ODOR STUDY COMMITTEE

NOVEMBER 28, 2007

PRESENTATION BY

DR. PATRICK G. HALBUR AND DR. JEFF ZIMMERMAN

COLLEGE OF VETERINARY MEDICINE IOWA STATE UNIVERSITY

A review of airborne transmission of pig diseases

College of Veterinary Medicine Iowa State University

ΔÎΔ

Iowa Legislature General Session Livestock Odor Study Committee November 28, 2007



Overview

- Airborne infectious diseases of pigs
- Research in airborne infectious diseases
- Current state of knowledge
 - Influenza and PRRS virus

Airborne infectious agents

- Domestic agents
 - Influenza viruses
 - PRRS virus (\$600M/yr)
 - Mycoplasma hyopneumoniae
 - Actinobacillus
 - pleuropneumoniae
 - Salmonella spp.
 - TGE virus

- "Exotic" agents
 - Foot-and-mouth disease virus
 - Hog cholera virus
 - African swine fever virus
 - Swine vesicular disease virus
- Pseudorabies virus

Aerosol research objectives:

- 1. Develop a body of knowledge
- 2. Predict events
- 3. Develop interventions



Aerosol research objectives:


- 1. Develop a body of knowledge
- 2. Predict events
- 3. Develop interventions





An aerosol example ...

- 1. Develop a body of knowledge
- 2. Predict events
- 3. Develop interventions



· March 1981, outbreaks of FMDV on 13 swine farms in Brittany and 1 in Normandy.



Knowledge → Prediction → *Intervention*


- · March 1981, outbreaks of FMDV on 13 swine farms in Brittany and 1 in Normandy.
- Prediction: transmission 186 miles to the coast of England.



Knowledge → Prediction → *Intervention*

- · March 1981, outbreaks of FMDV on 13 swine farms in Brittany and 1 in Normandy
- Prediction: transmission 186 miles to the coast of England.
- · Intervention:
 - Personnel in place
 - Detected infected cattle
 - Cattle removed
 - Catastrophe (\$) avoided






Knowledge → Prediction → *Intervention*


FMDV biology


- 1. How much FMDV shed by animals
- 2. Stability of airborne virus
- 3. Dose of virus required to infect animals



FMDV biology + meteorology

- Rate of virus decay under specific conditions.
 - Temperature
 - · Relative humidity
 - Cloud cover (UV)
- Virus dispersion
 - · Wind speed
 - · Wind direction





Current body of knowledge:
PRRS virus
Influenza virus



Body of knowledge

- General absence of knowledge on airborne transmission
- ... minimal understanding of the dynamics of virusladen aerosols ... ability to control and prevent virus spread is severely reduced, as was clearly demonstrated by SARS.

Morawska L. 2006. Droplet fate in indoor environments, or can we prevent the spread of infection? Indoor Air 16:335-347.

Body of knowledge

- General absence of knowledge on airborne transmission
- · Particularly true for livestock pathogens
- · No facts on which to base site placement



Body of knowledge

 Stärk KDC. 1999. The role of infectious aerosols in disease transmission in pigs. Veterinary Journal 158:164-181.

Disease	Field evidence	Indution from air	% RH for best survival	Menimal infection dose
Foot-and-mouth disease	500	98%	>60	2.6 log_ LD_
Swine vesicular disesse	no	909	>00	1
Anjeszky's disesse (Pseudorabies)	tera	909	55	4.5 log_TGD,
Influenca	tes	yes	3	3
Poecine respiratory and reproductive	pra	800	7	7
naothaw				
African mine fever	110	yes*	20-30	9
Classical meine fewy	0.0	80	2	7
Porcine respiratory coronavirus	tes	yes	7	7
Expostic pneumonia	term.	yes	<25 or >75+	7
Actinobacillosis	yes	yes	2	10%10° CFU/mL
Arthrophic shinits	100	yes	75	7

Body of knowledge

- · Systematic reviews in progress
 - PRRS virus (J. Hermann)
 - Influenza viruses (Dr. Annette O'Connor, Dr. Christa Irwin)




PRRSV - evidence of aerosol transmission • Experimental evidence (Wills et al., 1997) Programmental evidence group and grou

PRRSV - evidence of aerosol transmission

· Experimental evidence


aerosol 2/5 direct 5/5 indirect 3/5





PRRSV - evidence of aerosol transmission


- Experimental evidence
 - Transmission from inoculated to susceptible pigs over a distance of one meter in 1 of 2 attempts. (Torremorell et al., 1997)
 - ... 20 inches in 1 of 2 attempts. (Lager and Mengeling, 2000)
 - Transmission ... 2.5 m in infected animal air space, but not via exposure from exhaust fans over distances of 1 or 30 m. (Otake et al., 2002)
 - $-\,\dots$ one meter in 3 of 3 attempts. (Kristensen et al., 2002)

PRRSV - shedding by pigs - Hermann et al. 2007. Detection of respiratory pathogens in air samples from acutely infected pigs. Can J Vet Res (in press)









PRRSV - Summary

- · SHED at low levels
- STABLE at low temperature
- DOSE required to infect the next pig?
 In progress
- · Not enough data to determine site placement








Influenza - evidence of aerosol transmission

- · NO DATA equivalent in strength to PRRSV data
- · Transmission studies with guinea pigs, mice, ferrets
 - Lowen et al. 2007. Influenza virus transmission is dependent on RH and temperature. PLoS Pathog 3(10):e151
 - Lower temperature
 - Lower relative humidity




Influenza - shedding by pigs?


- NO DATA on respiratory aerosols from pigs
- Humans
 - Nasopharyngeal washes day 2 3 post infection
 - Up to 107 TCID₅₀ per ml
 - Big problems with indirect measures!!



Influenza - stability in the air

- · Limited data
- Mitchell et al. 1966. Decay of influenza A viruses of human and avian origin. Can J Comp Med Sci 32:544-546.





Influenza - infectious dose for pigs?

- · NO DATA on pigs
- Horses
 - 1 x 10² EID₅₀ lowest infectious dose
 Mumford et al. 1990. Experimental infection of ponies with equine influenza (H3N8) viruses by intranasal inoculation or exposure to aerosols. Equine Vet J 22:93-98
- Humans
 - 1 x 10^{0.6} to 1 x 10³ TCID₅₀
 Douglas RG, Influenza in man. 1975. In: The influenza viruses and influenza. New York: Academic Press. p. 375–447.
- Problems with extrapolation across species!!

Influenza - Summary

- · No data on respiratory SHEDDING by pigs
- Sparse data on STABILITY of airborne virus.
 Assumptions: better stability at lower RH and temps
- · No data on infectious DOSE
 - Highly infectious in equines and humans
- · NOT enough data to determine site placement








What do we need?

- Influenza
 - Shedding Stability Dose
 - IN PROGRESS: Koziel JA, Hoff SJ, Zimmerman JJ. Simultaneous treatment of odorants and pathogens emitted from CAFOs by advanced oxidation technologies. Binational Agricultural Research and Development Fund (BARD) - \$48,000.
- · PRRSV and influenza
 - "Baseline" field data



Baseline field data

- Objectives
 - Establish level and frequency of shedding
 - Estimate potential distance of spread
 - Evaluate effect of weather



Experimental design

- Monitor PRRSV and influenza on 25 growing pig sites
- 2. Site samples
 - 12 samples per site
 - oral fluid sampling (n = 5)
 - air sampling (n = 1)
- 3. Test samples for viruses
 - PCR
 - Determine virus relatedness among sites in cluster





Omni 3000™ air sampler

Baseline field data

BUDGET*	Г	
PRRSV PCV @ \$25 per sample		\$45,000
SIV PCR @ \$40 per sample		\$72,000
PRRSV sequencing (assume 6 per site) at \$150		\$22,500
SIV sequencing (assume 2 per site) at \$150		\$7,500
Two Omni™ 3000 @ \$25,000		\$50,000
Travel		\$10,000
Wages, salary, tuition, fringe (2 years)		\$70,000
TOTAL		\$277,000

^{*}Samples (25 sites x 12 samplings x 6 samples) = 1,800

Aerosol overview

- 1. Knowledge

 - Virus shedding by pigs
 Stability of airborne infectious virus
 Dose required to infect the next pig
- 2. Prediction
 - Where and when the virus will move
- 3. Intervention / Remediation
 - Filtration
 - Ultraviolet inactivation
 - Site placement

Thank you